

# Aptima® Multitest Swab Specimen Collection Kit

## Clinician collection procedure guide



**LABORATORY ALLIANCE**  
of Central New York, LLC

[laboratoryalliance.com](http://laboratoryalliance.com)

### Collection for vaginal, throat and rectal swab specimens



#### 1. Preparing to collect swab (regardless of source)


Partially open the swab package and remove the swab. Do not touch the soft tip or lay the swab down. **If the soft tip is touched, laid down, or dropped, discard and get a new Aptima Multitest Swab Specimen Collection Kit.** Hold the swab, placing thumb and forefinger in the middle of the shaft covering the black score line. Do not hold the shaft below the score line.

#### Swab specimen collection guide for:

- *Chlamydia trachomatis* (CT)
- *Neisseria gonorrhoeae* (NG)
- *Trichomonas vaginalis* (TV) for vaginal specimens only


#### 2. Collection of specimens:

##### vaginal




Carefully insert the swab into the vagina about 2 inches (5 cm) past the introitus and gently rotate the swab for 10 to 30 seconds. Make sure the swab touches the vaginal walls so that moisture is absorbed by the swab. Withdraw the swab without touching the skin.

##### throat



Carefully insert the swab into the throat ensuring contact with bilateral tonsils (if present) and the posterior pharyngeal wall, then withdraw the swab without touching the inside of the cheeks or tongue.


##### rectum



Carefully insert the swab into the rectum about 1-2 inches (3-5 cm) past the anal margin (the outside of the anus) and gently rotate the swab for 5 to 10 seconds. Withdraw the swab without touching the skin.



While holding the swab in hand, unscrew the tube cap. Do not spill the tube contents. If the tube contents are spilled, discard and replace with a new Aptima Multitest Swab Specimen Collection Kit. Immediately place the swab into the transport tube so the black score line is at the top of the tube. Align the score line with the top edge of the tube and carefully break the shaft. The swab will drop to the bottom of the tube. Discard the top portion of the swab shaft.



Tightly screw the cap onto the tube. When collecting multiple specimens from the same patient, the tube label provides a specimen source field for unique identification for the specimen location.



[hologic.com](http://hologic.com) • [diagnostic.solutions@hologic.com](mailto:diagnostic.solutions@hologic.com) • 888-484-4747

For more information, contact Laboratory Alliance of Central New York • 315-461-3008  
1304 Buckley Road, Suite 300, Syracuse, N.Y. 13212 • [laboratoryalliance.com](http://laboratoryalliance.com)